

The Dynamics, Structure and History of Galaxies: Workshop Summary

Joss Bland-Hawthorn

*Anglo-Australian Observatory, 167 Vimiera Road, Eastwood, NSW
 2122, Australia*

1. Introduction

As far as anyone knew, we were gathered at the resplendent Dunk Island on the Barrier Reef to celebrate Ken Freeman's 60th birthday. But what we got was a well presented cross-section of key topics in Galactic and extragalactic astronomy. Many of the speakers were former students of Ken's and the quality of the talks reflected this fact. If there was an overarching theme to the meeting, it reflected Ken's wish to understand "the chain of events that occurred during the formation of the Galaxy", to quote from his Annual Reviews article in 1987. The four day workshop was divided into four parts: galaxy structure and dynamics, haloes and environments, HI and dark matter, galaxy formation and evolution.

2. Galaxy Structure and Dynamics

Van der Kruit began the meeting with a bold statement: Ken Freeman's celebrated paper in 1970 remains the most highly cited in *Astrophysical Journal* of that year, and falls within the Top Ten of the most cited papers in galactic astronomy. He went on to review what we have learnt over the intervening years about galaxy disks, concluding that the central surface brightness μ_o given by "Freeman's law" – μ_o is approximately constant for spiral disks – is really an upper limit on μ_o . It is interesting to note that, in the original paper, Ken makes no statement that would constitute a law, although he does explore the theoretical consequences of constant μ_o . But the "Freeman disk" continues to play an important role in observational and theoretical studies.

The remarkable fact of the existence of the Tully-Fisher relation continues to puzzle. De Blok outlined the intimate connection of the Tully-Fisher conundrum with galaxy formation and evolution, a theme picked up by Bosma in his review of new HI data of late-type spirals. Bosma retraced the history of the dark matter problem and emphasized the importance of Ken's 1970 paper in stressing the mass discrepancy in galaxy dynamics, long before the spate of 'dark halo' theory papers starting in 1973. Throughout the 1970s, the observational case for dark matter became more compelling largely due to HI studies of spirals by Bosma, Roberts and others. From new observations of late-type spirals, Bosma finds the HI rotation curves are much better fit with an isothermal halo, rather than the NFW profile from CDM simulations.

Kormendy demonstrated the now-famous relation between the black hole mass and the depth of the potential well as measured by the velocity dispersion. There is a clear link between bulges and black holes which was probably established in the early universe during the major epoch of bulge growth. M33, which has no bulge, appears to have no black hole to an upper mass limit of 1000 solar masses. Sadler showed evidence for a radio-power/black hole mass relation which may allow us to track the evolution of black hole mass with redshift, something that is presently difficult to do for bulges.

Planetary nebulae are proving to be excellent probes of the outermost halo. Ford showed the kinematics of many hundreds of planetary nebulae out to 6 scale lengths in Cen A. The bisymmetric kinematic signature is strongly suggestive of a tumbling or triaxial potential. An obvious application would be to look for planetary nebulae in the faint stellar streams which are seen to encircle a number of nearby galaxies, including both M31 and the Galaxy.

Bureau, on behalf of the SAURON team, presented integral field results for a large sample of elliptical, lenticular and spiral galaxies divided between clusters and the field. Fully one third of these show evidence of nuclear substructure. Some show evidence for triaxiality, including nuclear bars as judged from Athanassoula's bar simulations. The group emphasizes caution in the analysis and derivation of black hole masses from longslit measurements.

Kalnajs entertained us with on-line simulations of Freeman bars and showed that they come in three flavours; slow, fast and pointy. Block presented a new physical classification of galaxies based on near-IR imaging. The method uses Elmegreen's proposal to estimate bar torque from the projected mass distribution. The scheme is based on the Fourier h_2 component of the IR light and the degree of arm winding. Block finds that "once you penetrate the mask," many multi-arm spirals are found to be grand design 2-arm spirals in the near IR. There appears to be a major overlap in bar torque between normal and barred spirals. This resonated with Gerhard who presented the crucial facts for a bar in the Galaxy. The predicted gas flow patterns reveal four arms – "four arms good, two arms bad" – which appear to agree with the tangent points for the spiral arms defined by HII regions. The COBE/DIRBE maps appear to require at least two spiral arms, but beyond that, not much can be deduced.

3. Haloes and environments

One of the most active research areas of astrophysics, especially so in Australia, concerns the chemical enrichment of the Galactic halo. From Norris' review, it was clear that we have barely begun to tap the information available from starlight. One of the most interesting results is the dramatic fall off in the number of stars with $[\text{Fe}/\text{H}] < -4$ with only a few currently known. A simple one-zone model of galactic chemical enrichment predicts that the number of stars should decline by a factor of ten for every decrease of 1 dex in $[\text{Fe}/\text{H}]$. Why this is not the case below $[\text{Fe}/\text{H}] < -4$ appears to be saying something important about the nature of the first stars in the universe.

Some of the heavy r- and s-process elements in metal-poor stars display an increasingly large scatter as $[\text{Fe}/\text{H}]$ decreases. This remarkable trend is thought to reflect incomplete mixing of the ISM in the early Galaxy. In his talk, Bland-

Hawthorn asked to what extent we could unravel disk formation from unique chemical signatures back to the last major epoch of dissipation.

Dwarf galaxies continue to present puzzles. Some like Sculptor and Fornax have low M/L values, while others like Sextans, Draco, Carina and Ursa Minor have huge overdensities of dark matter. It seems likely that some of the observed dwarfs have undergone dynamical stripping by the Galaxy. But Da Costa showed that these objects apparently cannot have contributed a large fraction of the halo stars: while the iron peak element abundances are much like field stars, the $[\alpha/\text{Fe}]$ ratio is substantially lower.

In the words of Mateo, the Local Group is a dangerous place for a dwarf. The tidal radius of a dwarf with mass m_7 (in units of $10^7 M_\odot$) at a distance r_{100} (in units of 100 kpc) from the centre of the Galaxy is roughly $r_d \sim r_{100} \sqrt[3]{m_7}$. The velocity dispersion of the dwarf is $\sigma_d \sim 6 \sqrt[3]{m_7} / \sqrt{r_{100}}$ (in units of km s^{-1}). We can approximate the timescale for dynamically stripping a dwarf galaxy as $\tau_d \sim 2\sqrt{3}(r_d/\sigma_d)$. A dwarf galaxy that ventures within 100 kpc of the two great rajahs of the Local Group can be stripped in just a few gigayears. In the coming decade, it is clear that detailed heavy element abundances (beyond those shown by Da Costa) will provide a clearer picture as to what fraction of the halo stars formed *in situ* rather than in satellite galaxies.

Future astrometric missions will no doubt find stellar streams associated with open clusters, globular clusters, and events unseen. Maybe all of the halo, much of the disk and some fraction of the bulge is made up of streams and substructure. Morrison spoke of the Spaghetti Survey which should be sensitive to such events out to a radius of 100 kpc. Clumping in the halo is also the main science goal of the YSTAR project, as presented by Byun.

De Zeeuw showed that the Gaia mission at the end of the decade will be able to measure the parallax and proper motions of 1.3 billion stars down to the level of micro-arcseconds. At these levels, the gravity field of Jupiter is detectable over the entire sky! Andrew Murray, author of an authoritative text on vectorial astrometry (“which dares not to venture below milli-arcsecond levels”), once remarked that it would take a very bright individual a lifetime to get to the bottom of all relativistic phenomena at micro-arcsecond levels. This of course provides a rich data set in its own right.

King presented his analysis, in collaboration with Jay Andersen, of Hubble Space Telescope data on M4 which allows for a near-perfect separation of the cluster main sequence from the field stars from the proper motions. The dispersion in the main sequence becomes almost immeasurable after the line-of-sight objects are rejected. This is reminiscent of recent work on open clusters which used the Hipparcos database to weed out objects not associated with the clusters. Quillen finds that the dispersion in $[\text{Fe}/\text{H}]$ is immeasurable once line-of-sight objects are rejected.

4. HI and Dark matter

While the “successes” of CDM on linear scales are trumpeted on a routine basis, this cannot be said for CDM predictions of the non-linear domain. This problem is particularly acute in regions of large density contrast. CDM simulations overpredict the cuspieness of galaxies, the dark matter fraction within the Solar

Circle, the random motions in the Local Supercluster, and the number satellites on the scale of the Local Group. Several speakers (e.g. Fall) noted that relaxing the cuspieness of CDM profiles makes everyone's life easier. But until this issue is resolved, interest in modified gravity laws will continue, even in the absence of an inertial framework. Sellwood showed the extent to which the MOND prescription can be made to explain a diverse array of data, particularly the fact that dark haloes appear to mimic the luminous components, e.g. squashed stellar systems have squashed dark haloes.

Sadler, Carignan and Putman presented some of the major results to derive from the southern sky HIPASS survey. It seems that the Ly-alpha absorption systems arise primarily in the low mass HI ellipticals are detected in HI in the HIPASS survey, and some of these structures stretch to vast distances. Gibson showed that most high-velocity clouds are likely to fall within 100 kpc of the Galaxy, although Putman finds that some fraction of these might extend further still. This, however, assumes that some compact clouds are physically associated with dark matter haloes. It is therefore noteworthy that Carignan claims evidence of HI associated with several dwarf spheroidals. In the Galaxy, Allen presented evidence for the association of HI with molecular gas which is being dissociated by the soft UV radiation field. This link can be seen over spatial scales which cover many orders of magnitude (1 pc to 10 kpc). Calzetti discussed how detailed observations of local galaxies can yield insight into the multiple processes associated with the evolution of starbursts.

The detailed connection of baryons to dark matter continues to elude. We should prepare ourselves for the possibility that there is more to the dark matter universe than we might suppose. CDM appears to predict that the dark matter universe is largely uninteresting. But the galaxies we know and love are the light cusps within the dark matter potential, much as cD galaxies are to galaxy clusters. Dark matter properties may well set up the visible universe to the extent that local dark matter properties correlate with Hubble type and luminosity.

5. Galaxy formation and evolution

Fall showed that the salient features of the Fall & Efstathiou model of disk formation survive within the CDM hierarchy. But, as Weil et al. (1998, MNRAS, 300, 773) point out, this does require that most of the gas in small systems stays hot long enough to settle to the disk when most of the accretion activity has passed. Fall and Silk both emphasized the importance of viscosity and its relation to star-formation timescales. In the Lin & Pringle picture, this viscosity is maintained by the disk-halo interaction and spiral arms.

Jerjen demonstrated that the Surface Brightness Fluctuation method can be used to determine distances to dwarf elliptical galaxies. This then opens the way for mapping in a simple and effective way the detailed distribution of galaxies within ~ 10 Mpc. This may allow us to understand how large scale structures within this volume, such as the Supergalactic Plane, arise. Student *numero uno* Illingworth showed that the universe at $z \sim 1$ looks much like the universe today. He emphasized that since most of the baryons reside in ellipticals and bulges today, these largely define galaxy evolution.

6. Conclusions

We are coming into a new era of Galactic investigation where it is now possible to identify fossil evidence of ancient events. In the Galactic halo, we might label these the Helmi event, the ω Cen event, the Sgr dwarf event and the “thick disk” event (i.e. the event which gave rise to the thick disk). Some of these halo objects may even be related.

In fact, there are likely to be many signatures of the galaxy formation process, not only substructure in the outer halo (Helmi, Morrison). Some of these, like the existence of truncation radii (van der Kruit) and the distribution of specific angular momentum between the Galactic components (Fall), may have been set down long ago. In all likelihood, we have yet to recognize many important signals, i.e. fossil signatures, of past events.

One such signature may be the existence of superthin disk galaxies, examples of which were shown in several talks. It is interesting that superthin disk galaxies exist at all. These may be telling us that, for the most part, disk formation is highly quiescent. It simply putters along over 10 Gyr or so. It is tempting to suggest that the Madau plot, which purports to monitor the star formation history over cosmic time, is largely dominated by star formation in merger events. In this picture, all disk thickening is due to dynamical events like accretion.

To come back to Ken’s original wish, it is becoming increasingly clear that “to understand the chain of events that occurred during the formation of the Galaxy”, the astronomical community *must embrace the wealth of near-field information with the same relish that it embraces the far field universe*. There are many compelling reasons to support this statement. In particular, two key events in the early universe are the epoch when dark matter haloes first became established (roughly 14 Gyr ago), and the epoch when disks first established themselves (10–12 Gyr ago). The world-model parameters ($\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$) place both of these events in the redshift window which defines contemporary observational cosmology ($z \approx 1 - 6$) at optical/IR wavelengths.

Many of the luminous objects we observe at high redshift are chemically and dynamically evolved as shown by solar abundance levels (the dynamical timescales are short in the cores of galaxies). To understand galaxy evolution, we would like information from the outer parts of high-redshift galaxies, which is exceedingly hard to obtain. This information can be extracted from near-field studies although now the systems are viewed at a much later stage in their evolution. In any event, near-field and far-field data are entirely complementary, and a successful theory of galaxy formation will need to be firmly anchored to both extremes.